




*EIC Comprehensive  
Chromodynamics  
Experiment*



# ECCE Physics performance

Carlos Muñoz Camacho


IJCLab-Orsay, CRNS/IN2P3 (France)

# Key science questions that the EIC will address




## Origin of spin


- How does the spin-1/2 of the nucleon arise from the spin of quarks, gluons and their orbital angular momenta?




## Origin of mass



- How do massless gluons make up for most of the nucleon mass?



## Gluons in nuclei



- Does gluon density saturate at high energy giving rise to a new regime of matter?

- ECCE was designed to address all these questions
- Detailed analyses based on full Geant4 simulations demonstrate that **ECCE can deliver on the science outlined in the EIC White Paper and the NAS report**
- All studies are documented in dedicated analysis notes: <https://www.ecce-eic.org/ecce-internal-notes> (pwd: ECCEprop)

# Origin of nucleon spin: physics requirements



## Physics measurements:

- Inclusive Deep Inelastic Scattering (DIS) measurements

- Double spin asymmetry  $A_1(x)$

Good scattered electron identification and resolution

High resolution homogeneous ECALs in both backwards endcap and barrel

Good momentum resolution with hybrid AI-optimized tracker

- Semi-inclusive DIS measurements


- Collins & Sivers asymmetries

Low  $p_T$  acceptance down to 100 MeV for  $\pi$


1.4T field

$3\sigma$  h-PID up to 8 GeV/c (backwards) & up to 6 GeV/c (central) & up to 50 GeV/c (forward)

High performance Cherekov PID systems for large momenta complemented by TOF systems for low-p PID




# Origin of nucleon spin: physics performance




Full Geant4 simulation

**Systematic uncertainties:**  
difference between  
reconstructed & truth  
kinematics



Transverse spin asymmetry as a function of hadron momentum  $z$




Inclusive double spin asymmetry  $A_1(x)$



Hermeticity  $-3.5 < \eta < 3.5$  and low momenta  $\pi$  acceptance



Impact on  $u$  and  $d$   
transversity distributions



Impact on favored & disfavored  
Collins fragmentation functions



Accuracy compatible  
with projections in  
the YR

# Origin of the mass: requirements



## Physics measurements:

- Exclusive reactions with very forward particle detection/tagging
- High acceptance up to very small  $p_T$
- Excellent  $p_T$  resolution
- Precise timing (correction crab cavity rotation)
- High precision tracking and timing (AC-LGADs) in all B0, Roman Pots & Off-Momentum detectors
- Zero-Degree Calorimeter (ZDC) design as developed during the YR
- Lead-tungsten calorimeter in B0 magnet to measure physics beyond the WP & NAS report (eg. u-channel DVCS)




# Origin of mass: physics performance (1)



## 3D imaging of quarks and gluons





- Full simulation of several exclusive channels (DVCS, exclusive  $J/\psi$ ...)
- Beam effects (cross-angle & beam divergence) included
- Large and continuous coverage in  $t$  ( $\sim p_T^{-2}$ ) up to very small values of  $p_T$

Exclusive  $J/\psi$  production



ECCE can deliver the physics outlined in the WP and NAS report

t-dependence of DVCS cross section





# Origin of mass: physics performance (2)




- Threshold  $J/\Psi$  production as a function of  $Q^2$  is sensitive to the trace anomaly contribution to the proton mass
- Meson structure measurements (structure function & form factor) probe the hadron mass generation through chiral symmetry breaking

Tracking capabilities of ECCE enable  $J/\Psi$  over wide range in  $t$  and  $Q^2$



Far-forward detectors are key for reconstructing mesons at very  $\eta$




# Gluons in nuclei: physics requirements





## Physics measurements:

- SIDIS heavy flavor production
  - Good vertex resolution for open heavy flavor reconstruction
  - AI-optimized hybrid Si tracker
- Dijet measurements
  - Good hadron calorimetry
  - Reuse of existing hadronic calorimeters
- Diffractive processes off nuclei
  - Excellent backwards EMCal
  - Choice of full  $PbWO_4$  crystal calorimeter




Distance of closest approach within YR requirements



Reconstruction for track+cluster jets exceeds YR requirements

Excellent EMCAL energy resolution






# Gluons in nuclei: physics performance (1)




## Gluon saturation


- Early nPDF measurements will probe gluon saturation regime (by comparing to DGALP evolution and by using different nuclei)
- High resolution backwards EMCAL allows to distinguish change in slope in diffractive production
- Jet reconstruction sensitive to saturation effect in eg. dihadron correlations



Diffractive  $J/\psi$  production off Pb



Dihadron correlation from dijet production in ep & eAu






# Glueons in nuclei: physics performance (2)



## Nuclear matter and hadronization

- SIDIS in eA is an excellent process to understand hadronization
- Heavy flavor (HF) production provides a clean probe of gluon dynamics in nucleons and nuclei
- Comparison of HF production in ep & eA ( $R_{eA}$ ) proves the hadronization process in vacuum and in a cold nuclear medium

Nuclear modification factor in ep vs eAu for  $D^0$  ( $\bar{D}^0$ ) as a function of  $z_h$



- ✓ Tracking reconstruction of ECCE provides the necessary discriminating power between different model predictions of hadronization
- ✓ Acceptance for low momentum pions significantly increases statistical uncertainties


# Science beyond the NAS report (1)




## Precision electroweak and BSM physics

- The ECCE consortium is interested in parity-violating (PV) asymmetries and charged lepton flavor violating (CLFV) processes to search for physics BSM
- Using the DIS reconstruction capabilities and  $100 \text{ fb}^{-1}$  integrated luminosity, ECCE will set stringent limits in BSM physics


**PVDIS asymmetries:** Expected limits on 2 sets of Wilson coefficients associated with BSM degrees of freedom in the SMEFT



**Leptoquarks:** limits on contact interaction terms based on  $ep \rightarrow \tau X$  cross section



**Resolution in  $t$  decay length: 119 mm**




# Science beyond the NAS report (2)



## XYZ Spectroscopy

- Photoproduction of “XYZ” meson states probes underlying dynamics and allow determining their quantum number
- Detection of *low energy pions* is crucial while providing *good invariant mass resolution* :  
1.4T field is optimal for spectroscopy





- Access to nucleon *Transition Distribution Amplitudes* (TDA): light-cone matrix elements complementary to GPDs
- Bethe-Heitler is suppressed in the *u*-channel, but  $\pi^0$  background suppression is needed via an EMCAL at very forward rapidity (B0 magnet)

- Reconstructed invariant mass for 3 simulated states:  
 $\chi_{c1}(3872)$ ,  $Y(4260)$  and  $\Psi(2s)$
- 30 MeV resolution achieved with ECCE

Low- $Q^2$  tagger (far-backwards region) is crucial for this measurement

## *u*-channel DVCS



# Conclusion



- ECCE design was driven by the different physics measurements required to address the full set of EIC science
  - Each subsystem technology was chosen to address specific physics requirements
- ECCE can deliver on *all* these physics measurements and we have demonstrate this through full Geant4 simulations
  - ECCE meets (or exceeds) the detector requirements outlined in the Yellow Report
  - ECCE can address all of the physics topics listed in the EIC White Paper and the NAS report
  - ECCE physics performance compatible with projections from the Yellow Report exercise
- ECCE can also address several exciting physics topics beyond the WP and the NAS report
- All studies thoroughly documented in 15 physics analysis notes (+10 detector notes):  
<https://www.ecce-eic.org/ecce-internal-notes> (pwd: ECCEprop)