




arXiv 1212.1701.v3  
Eur. Phys. J. A52, 9 (2016)

# Electron Ion Collider: The next QCD frontier

*Understanding the **Glue** that Binds Us All*


This talk is based on the work of a large number of scientists, excited about the EIC science and involved in the EIC project, now organized as the EIC Users Group








Abhay Deshpande

# Why an Electron Ion Collider











- **Interactions and structure are mixed up in nuclear matter:** Nuclear matter is made of quarks that are bound by gluons that also bind themselves. Unlike with the more familiar atomic and molecular matter, the **interactions and structures are inextricably mixed up**, and the **observed properties** of nucleons and nuclei, such as mass & spin, **emerge** out of this complex system.
- **Gaining understanding of this dynamic matter → transformational:** Gaining **detailed knowledge** of this astonishing dynamical system at the heart of our world **could be transformational**, perhaps in an even **more dramatic way** than how the understanding of the atomic and molecular structure of matter led to new frontiers, new sciences and new technologies.
- **The Electron Ion Collider is the right tool:** A new US-based facility, high-energy, high-luminosity Electron Ion collider (EIC), capable of a versatile range of beam energies, polarizations, and species, is **required to precisely image the quarks and gluons and their interactions**, to explore the **new QCD frontier of strong color fields** in nuclei – to *understand* how matter at its most fundamental level is made.


















# EIC: A Portal to a New Frontier

Dynamical System	Fundamental Knowns	Unknowns	Breakthrough Structure Probes (Date)	New Sciences, New Frontiers
<p>Solids</p> 	<p>Electromagnetism Atoms</p> 	<p>Structure</p> 	<p>X-ray Diffraction (~1920)</p> 	<p>Solid state physics Molecular biology</p> 

# EIC: A Portal to a New Frontier



Dynamical System	Fundamental Knowns	Unknowns	Breakthrough Structure Probes (Date)	New Sciences, New Frontiers
<p>Solids</p> 	<p>Electromagnetism Atoms</p> 	<p>Structure</p>  <p>1801</p>	<p>X-ray Diffraction (~1920)</p> 	<p>Solid state physics Molecular biology</p>  <p>DNA</p>
<p>Universe</p> 	<p>General Relativity Standard Model</p> 	<p>Quantum Gravity, Dark matter, Dark energy. Structure</p>  <p>CMB 1965</p>	<p>Large Scale Surveys CMB Probes (~2000)</p> 	<p>Precision Observational Cosmology</p> 

# EIC: A Portal to a New Frontier

Dynamical System	Fundamental Knowns	Unknowns	Breakthrough Structure Probes (Date)	New Sciences, New Frontiers
<p>Solids</p> 	<p>Electromagnetism Atoms</p> 	<p>Structure</p>  <p>1801</p>	<p>X-ray Diffraction (~1920)</p> 	<p>Solid state physics Molecular biology</p>  <p>DNA</p>
<p>Universe</p> 	<p>General Relativity Standard Model</p> 	<p>Quantum Gravity, Dark matter, Dark energy. Structure</p>  <p>CMB 1965</p>	<p>Large Scale Surveys CMB Probes (~2000)</p> 	<p>Precision Observational Cosmology</p> 
<p>Nuclei and Nucleons</p> 	<p>Perturbative QCD Quarks and Gluons</p> $\mathcal{L}_{\text{QCD}} = \bar{\psi}(i\partial - g\mathbf{A})\psi - \frac{1}{2}\text{tr} F_{\mu\nu}F^{\mu\nu}$ 	<p>Non-perturbative QCD Structure</p>  <p>2017</p>	<p>Electron-Ion Collider (2025+)</p> 	<p>Structure &amp; Dynamics in QCD</p> 


# QCD Landscape explored by EIC

QCD at high resolution ( $Q^2$ ) — weakly correlated quarks and gluons are well-described



# QCD Landscape explored by EIC


QCD at high resolution ( $Q^2$ ) —weakly correlated quarks and gluons are well-described



Strong QCD dynamics creates many-body correlations between quarks and gluons  
 → hadron structure emerges


EIC systematically explores correlations in this region.

An exciting opportunity: Observation by EIC of a new regime in QCD of weakly coupled high density matter



# Non-linear Structure of QCD: Fundamental Consequences

- Quark (Color) confinement:
  - Consequence of nonlinear **gluon self-interactions**
  - Unique property of the strong interaction
- Strong **Quark-Gluon** Interactions:
  - **Confined motion** of quarks and gluons – Transverse Momentum Dependent Parton Distributions (TMDs)
  - **Confined spatial correlations** of quark and gluon distributions – Generalized Parton Distributions (GPDs)
- Ultra-dense color (**gluon**) fields:
  - Is there a universal many-body structure due to ultra-dense color fields at the core of **all** hadrons and nuclei?






# Emergent Dynamics in QCD

*Without gluons, there would be no nucleons,  
no atomic nuclei... no visible world!*



- Massless gluons & almost massless quarks, *through their interactions*, generate most of the mass of the nucleons
- Gluons carry ~50% of the proton's momentum, a significant fraction of the nucleon's spin, and are essential for the dynamics of confined partons
- Properties of hadrons are **emergent phenomena** resulting not only from the equation of motion but are also inextricably tied to the properties of the QCD vacuum. Striking examples besides confinement are spontaneous symmetry breaking and anomalies
- The nucleon-nucleon forces emerge from quark-gluon interactions: how this happens remains a mystery

**Experimental insight and guidance crucial for complete understanding of *how* hadrons & nuclei emerge from quarks and gluons**



# A new facility is needed to investigate, with precision, the dynamics of gluons & sea quarks and their role in the structure of visible matter



How are the sea quarks and gluons, and their spins, distributed in space and momentum inside the nucleon?  
How do the nucleon properties emerge from them and their interactions?



# A new facility is needed to investigate, with precision, the dynamics of gluons & sea quarks and their role in the structure of visible matter

How are the sea quarks and gluons, and their spins, distributed in space and momentum inside the nucleon?


How do the nucleon properties emerge from them and their interactions?



How do color-charged quarks and gluons, and colorless jets, interact with a nuclear medium?

How do the confined hadronic states emerge from these quarks and gluons?


How do the quark-gluon interactions create nuclear binding?



# A new facility is needed to investigate, with precision, the dynamics of gluons & sea quarks and their role in the structure of visible matter

How are the sea quarks and gluons, and their spins, distributed in space and momentum inside the nucleon?


How do the nucleon properties emerge from them and their interactions?



How do color-charged quarks and gluons, and colorless jets, interact with a nuclear medium?


How do the confined hadronic states emerge from these quarks and gluons?

How do the quark-gluon interactions create nuclear binding?




How does a dense nuclear environment affect the quarks and gluons, their correlations, and their interactions?

What happens to the gluon density in nuclei? Does it saturate at high energy, giving rise to a gluonic matter with universal properties in all nuclei, even the proton?





gluon emission




?

gluon recombination




# Uniqueness of EIC among all DIS Facilities




All DIS facilities in the world.

However,  
if we ask for:




# Uniqueness of EIC among all DIS Facilities




All DIS facilities in the world.

However,  
if we ask for:

- high luminosity & wide reach in  $\sqrt{s}$



# Uniqueness of EIC among all DIS Facilities




All DIS facilities in the world.


However,  
if we ask for:

- high luminosity & wide reach in  $\sqrt{s}$
- polarized lepton & hadron beams
- nuclear beams

**EIC stands out as  
unique facility ...**




# Uniqueness of EIC among all DIS Facilities






# The world's first polarized electron-proton collider

## Polarized proton as a laboratory for QCD



- How are the sea quarks and gluons, and their spins, *distributed in space and momentum* inside the nucleon?
- How do the *nucleon properties emerge* from them and their interactions?




# What does a proton look like with increasing energy?

**One** of several possible scenarios: a pion cloud model

A parton core in the proton gets increasingly surrounded by a meson cloud with decreasing  $x$

→ large impact on gluon and sea-quark observables



## What do we expect to see:


- $q\bar{q}$  pairs (sea quarks) generated at small(ish)- $x$  are predicted to be unpolarized
- gluons generated from sea quarks are unpolarized

→ needed:


- high precision measurement of flavor separated polarized quark and gluon distributions as functions of  $x$
- high precision spatial imaging: **Gluon radius ~ sea-quark radius ?**

## What happens in the gluon dominated small- $x$ regime?

- possible scenario: lumpy glue




**EIC will explore the dynamical spatial structure of hadrons**



# 3-Dimensional Imaging Quarks and Gluons


## Wigner functions $W(x, b_T, k_T)$

offer unprecedented insight into confinement and chiral symmetry breaking.



Spin-dependent 3D **momentum space** images from semi-inclusive scattering  
→ **TMDs**


Spin-dependent 2D **coordinate space** (transverse) + 1D (longitudinal momentum) images from exclusive scattering  
→ **GPDs**



# 3-Dimensional Imaging Quarks and Gluons

## Wigner functions $W(x, b_T, k_T)$

offer unprecedented insight into confinement and chiral symmetry breaking.




Spin-dependent 3D **momentum space** images from semi-inclusive scattering  
→ **TMDs**

Spin-dependent 2D **coordinate space** (transverse) + 1D (longitudinal momentum) images from exclusive scattering  
→ **GPDs**

Position and momentum → Orbital motion of quarks and gluons


Recent theoretical work indicates possible direct access to gluon Wigner function through diffractive di-jet measurements at an EIC



# 2+1 D partonic image of the proton with the EIC

Spin-dependent 3D **momentum space**  
images from semi-inclusive scattering

Spin-dependent 2D **coordinate space**  
(transverse) + 1D (longitudinal momentum)  
images from exclusive scattering





# 2+1 D partonic image of the proton with the EIC

Spin-dependent 3D momentum space images from semi-inclusive scattering

Spin-dependent 2D coordinate space (transverse) + 1D (longitudinal momentum) images from exclusive scattering

## Transverse Momentum Distributions





# 2+1 D partonic image of the proton with the EIC

Spin-dependent 3D momentum space images from semi-inclusive scattering

Spin-dependent 2D coordinate space (transverse) + 1D (longitudinal momentum) images from exclusive scattering


## Transverse Momentum Distributions



# 2+1 D partonic image of the proton with the EIC

Spin-dependent 3D momentum space images from semi-inclusive scattering


## Transverse Momentum Distributions



Spin-dependent 2D coordinate space (transverse) + 1D (longitudinal momentum) images from exclusive scattering

## Transverse Position Distributions

sea-quarks  
unpolarized      polarized






# 2+1 D partonic image of the proton with the EIC



Spin-dependent 3D momentum space images from semi-inclusive scattering

## Transverse Momentum Distributions




Spin-dependent 2D coordinate space (transverse) + 1D (longitudinal momentum) images from exclusive scattering

## Transverse Position Distributions




# Understanding Nucleon Spin





“Helicity sum rule”

$$\frac{1}{2}\hbar = \underbrace{\frac{1}{2}\Delta\Sigma}_{\text{quark contribution}} + \underbrace{\Delta G}_{\text{gluon contribution}} + \underbrace{\sum_q L_q^z + L_g^z}_{\text{orbital angular momentum}}$$


EIC projected measurements:  
 precise determination of polarized PDFs of quark sea and gluons → precision  $\Delta G$  and  $\Delta\Sigma$   
 → A clear idea of the magnitude of  $\sum L_q + L_g$



DIS + SIDIS with 90% C.L. band  
 DIS + SIDIS + RHIC with 90% C.L. band  
 RHIC projection including 500 GeV data  
 EIC projection  $\sqrt{s} = 78$  GeV





# Understanding Nucleon Spin



“Helicity sum rule”

$$\frac{1}{2}\hbar = \underbrace{\frac{1}{2}\Delta\Sigma}_{\text{quark contribution}} + \underbrace{\Delta G}_{\text{gluon contribution}} + \underbrace{\sum_q L_q^z + L_g^z}_{\text{orbital angular momentum}}$$

EIC projected measurements:  
 precise determination of polarized PDFs of quark sea and gluons → precision  $\Delta G$  and  $\Delta\Sigma$   
 → A clear idea of the magnitude of  $\sum L_q + L_g$




## Spin and Lattice: Recent Activities


❑ **Gluon’s spin contribution on Lattice:  $S_G = 0.5(0.1)$**   
 Yi-Bo Yang et al. PRL **118**, 102001 (2017)

❑  **$J_q$  calculated on Lattice QCD:**  

 QCD Collaboration, PRD91, 014505,



# Understanding Nucleon Mass



Relativistic motion

$\chi$  Symmetry Breaking

Quantum fluctuation

$$M = E_q + E_g + \chi m_q + T_g$$

Quark Energy

Gluon Energy


Quark Mass

Trace Anomaly



“... The vast majority of the nucleon’s mass is due to quantum fluctuations of quark-antiquark pairs, the gluons, and the energy associated with quarks moving around at close to the speed of light. ...”

*The 2015 Long Range Plan for Nuclear Science*

## □ Preliminary Lattice QCD results:




# Understanding Nucleon Mass



“... The vast majority of the nucleon’s mass is due to quantum fluctuations of quark-antiquark pairs, the gluons, and the energy associated with quarks moving around at close to the speed of light. ...”

*The 2015 Long Range Plan for Nuclear Science*


## □ Preliminary Lattice QCD results:



## □ EIC’s expected contribution in:

- ◇ Trace anomaly:

*Upsilon production near the threshold*




- ◇ Quark-gluon energy:

$\propto$  quark-gluon momentum fractions




*In nucleon with DIS and SIDIS*

*In pions and kaons with Sullivan process*




# The world's first electron-nucleus collider

## The Nucleus as a laboratory for QCD



- How do color-charged quarks and gluons, and colorless jets, *interact with a nuclear medium*?
- How do *the confined hadronic states emerge* from these quarks and gluons?
- How does the quark-gluon interaction *create nuclear binding*?




# Emergence of Hadrons from Partons

## Nucleus as a Femtometer sized analyzer

Unprecedented  $\nu$ , the virtual photon energy range @ EIC : precision & control


$$\nu = \frac{Q^2}{2mx}$$



Control of  $\nu$  by selecting kinematics;  
Also under control the nuclear size.

Colored quark emerges as color neutral hadron → What is the impact of colored media on confinement?

Energy loss by light vs. heavy quarks:



Identify light vs. charm hadrons in e-A:  
Understand energy loss of light vs. heavy quarks in cold nuclear matter.  
Provides insight into energy loss in the Quark-Gluon Plasma

**DIS at collider energies enables control of parton/event kinematics**


# Jets: A Window to Partons

- Jet: transition process from a parton to hadrons and as such fundamentally encodes hadronization and dynamic confinement.
- Jets are golden tools to study quarks and gluons at RHIC and LHC
- Jets probe the interaction with the medium using the well understood jet shower evolution to extract the space-time dynamics of hadronization.

## Jets in eA Collisions

- Jet showers, their correlations and attenuation shed light on hadronization and dynamical nature of confinement in extended colored media
- Determine the transport properties in cold QCD medium
- Gluon distribution from dijets

**Many opportunities for jet physics in polarized ep**







# Short range correlations & physics with light nuclei

Exciting area of interest:


$$e + D \rightarrow e' + p + n + J/\Psi$$

Exclusive measurements of tagged (polarized) protons and neutrons in coincidence with vector mesons probe the short-range quark-gluon nature of nuclear forces





Tag the recoil proton  $\rightarrow$  Study the neutron's spin structure function.

Other possibilities: Polarized EMC effect with polarized light nuclei




# The world's first electron-nucleus collider


## The Nucleus as a laboratory for QCD



- How does a *dense nuclear environment* affect the quarks and gluons, their correlations, and their interactions?
- What happens to the *gluon density in nuclei*? Does it *saturate at high energy*, giving rise to a gluonic matter with *universal properties* in all nuclei, even the proton?



# EIC: impact on the knowledge of 1D Nuclear PDFs




## Ratio of Parton Distribution Functions of Pb over Proton:

- ❖ Without EIC, large uncertainties in **nuclear sea quarks and gluons**  
→ With EIC **significantly reduced uncertainties**
- ❖ Complementary to RHIC and LHC pA data. Provides information on initial state for heavy ion collisions.
- ❖ **Does the nucleus behave like a proton at low-x? → such color correlations relevant to the understanding of astronomical objects**




# Gluon saturation at low-x




## What tames the low-x rise?

- New evolution equations at low x & moderate  $Q^2$
- **Saturation Scale  $Q_s(x)$**  where gluon emission and recombination become comparable




gluon emission



=


gluon recombination



At  $Q_s$




# Gluon saturation at low-x




## What tames the low-x rise?

- New evolution equations at low x & moderate  $Q^2$
- **Saturation Scale  $Q_s(x)$**  where gluon emission and recombination become comparable




gluon emission



=

gluon recombination



At  $Q_s$

First observation of gluon recombination effects in nuclei:

→ leading to a **collective gluonic system**

First observation of gluon recombination in different nuclei

→

Is this a **universal property**?



What is the new effective theory in this regime?



# How to explore/study this new phase of matter?



(multi-TeV) e-p collider (LHeC) OR [a \(multi-10s GeV\) e-A collider](#)

## Advantage of nucleus →



$$(Q_s^A)^2 \approx c Q_0^2 \left[ \frac{A}{x} \right]^{1/3}$$




$$L \sim (2m_N x)^{-1} > 2 R_A \sim A^{1/3}$$



# How to explore/study this new phase of matter?

(multi-TeV) e-p collider (LHeC) OR [a \(multi-10s GeV\) e-A collider](#)



## Advantage of nucleus →



# How to explore/study this new phase of matter?

(multi-TeV) e-p collider (LHeC) OR [a \(multi-10s GeV\) e-A collider](#)

## Advantage of nucleus →



Enhancement of  $Q_s$  with  $A$ :  
Saturation regime reached at significantly lower energy  
(read: “cost”) in nuclei




# Diffraction for the 21<sup>st</sup> Century


Diffraction cross-sections have strong discovery potential:

High sensitivity to gluon density in linear regime:  $\sigma \sim [g(x, Q^2)]^2$


Dramatic changes in cross-sections with onset of non-linear strong color fields





Extracting the gluon distribution  $\rho(b_T)$  of nuclei via Fourier transformation of  $d\sigma/dt$  in diffractive  $J/\psi$  production




Probing gluon saturation through measuring  $\sigma_{diff}/\sigma_{tot}$



Probing  $Q^2$  dependence of gluon saturation in diffractive vector meson production




# EIC Science and Connections



# Connections to other areas of physics


- Explorations of the stringy dynamics of hadrons led to the string theory of Gravity. A weakly coupled regime of 10-d **gravity** is conjectured to be dual to strongly coupled 4-d QCD-like theory. *Further profound connections may emerge from deeper investigations of the QCD landscape.*
- The dynamics of strongly coupled **cold atom gases** and QCD (non-Abelian gauge fields but also strong nuclear fields) show strikingly common features. Cold atom scientists are actively engaged in engineering cold atoms simulators of gauge field mechanism.
- Strong connections have emerged between studies of **strongly correlated condensed matter systems** and QCD: *topological effects arising from chiral anomaly*
- **Strong field QED** explores the breakdown of the QED vacuum and its nonlinear optical response in  $e^+e^-$  pair creation. *Reaching this regime is a major goal in developing high powered lasers.*




# Collider Realization & US leadership in Accelerator Science

Collider requirements: High luminosity & energy, variable CM energy, all- A nuclear beams, polarization in e- and light ions of the EIC poses unique & attractive challenges to the accelerator physicists


→ *EIC demands frontier ideas and technologies in accelerator physics*




# JLEIC Realization




- Use existing CEBAF for polarized electron injector
- Figure 8 Layout: Optimized for high ion beam polarization → polarized deuterons
- Energy Range:  $\sqrt{s}$  : 20 to 65 - 140 GeV (magnet technology choice)
- Fully integrated detector/IR
- JLEIC achieves initial high luminosity, with technology choice determining initial and upgraded energy reach




# eRHIC Realization



- Use existing RHIC
  - Up to 275 GeV protons
  - Existing: tunnel, detector halls & hadron injector complex
- Add 18 GeV electron accelerator in the same tunnel
  - Use either high intensity Electron Storage Ring or Energy Recovery Linac
- Achieve high luminosity, high energy e-p/A collisions with full acceptance detector
- Luminosity and/or energy staging possible




# EIC Users Group



“....This facility can lead to the convergence of the present world-leading QCD programs at CEBAF and RHIC in a single facility.....”

## EIC Users Group

- Established, enthusiastic and active
- New physics ideas initiated with new influx of people....




# The EIC Users Group: EICUG.ORG

(no students included as of yet)

**685 collaborators, 28 countries, 151 institutions... (April, 2017)**

Map of institution's locations






# The EIC Users Group: EICUG.ORG

(no students included as of yet)

**685 collaborators, 28 countries, 151 institutions... (April, 2017)**

Map of institution's locations



**The EIC Users Meeting at Stony Brook, June 2014:**

→ <http://skipper.physics.sunysb.edu/~eicug/meeting1/SBU.html>

**The EIC UG Meeting at University of Berkeley, January 6-9, 2016**

<http://skipper.physics.sunysb.edu/~eicug/meeting2/UCB2016.html>

**Recent EICUG Argonne National Laboratory July 7-10, 2016**

<http://eic2016.phy.anl.gov>

**Remote/Internet: meeting: March 16<sup>th</sup> : For NAS Review preparation**


***Next meeting:***

***July 18-22, 2017 Trieste, Italy***



# New Users → New Physics → Lots of activities


- Jet studies at the EIC:
  - Systematic investigations of general issues in jet-finding at an EIC
  - Understanding of “micro-jets” – jets with only few hadrons
  - Understanding the jet structure modifications in nuclei vs. protons
  - Energy loss in cold QCD matter (Nuclei) vs. hot QCD matter at RHIC and LHC
- Precision measurements of the “initial state” for collisions leading to the QGP being studied at RHIC and LHC
- Precision PDF measurements in proton, neutron & photons at the EIC:
  - Study the free neutron PDFs through tagging and on-shell extrapolation
  - Study the gluon PDFs at large Bjorken  $x$  through evolution and open-charm production
  - Study of gluons TMDs
  - Study the potential impact on Higgs studies in the High-Luminosity LHC era
  - Study the impact of TMDs @ EIC on W-production at the LHC
  - Polarized and unpolarized photon PDFs
- Measurements of PDFs in pions and kaons through the Sullivan process
  - Theoretical studies of the equivalence of near-off-shell and on-shell pions and kaons
  - Study the extraction of, and expected differences of, quark and gluon PDFs in pions, kaons and nucleons, and the relation to their physical masses
- Nucleon structure with electroweak probes, and precision BSM physics (i.e.  $\text{Sin}^2\Theta_W$ )
- Heavy quark & quarkonia production with 100-1000 times HERA luminosity
- In view of new discoveries of multi-quark XYZ states: what could EIC contribute?



# New Users → New Physics → Lots of activities



**POETIC VI**  
6th International Conference on  
Physics Opportunities at an Electron-Ion Collider  
7-11 September 2016  
Ecole Polytechnique, Palaiseau, France  
<http://poetic6.scienceconf.org/>




**INSTITUTE FOR NUCLEAR THEORY**  
Home | Contact | Search | Site Map



International Advisory Committee  
H. Avancini, M. Bruneau, C. B. Dover, M. Galassi, M. L. Hellmich, M. J. Leitch, K. Li, M. M. S. Maciucă, M. S. Zisman, L. S. Zeng, A. Zang, T. Ziegler, R. Zito




**The Proton Mass**  
At the heart of most visible matter.  
Temple University, March 28-29, 2016

Speakers:  
Sean Bladys (SLAC), Xiaohong Di (Maryland), Dima Khazanchi (Brookhaven), Keith Hill (University of Kentucky), David Roberts (Edinburgh), Craig Roberts (ANL), Martin Senger (University of Washington), Stephen Seymour (JLab), George Sorensen (Brookhaven)

Moderator: Alfred Mueller (Columbia)

Local Organizers: Zainuddin Mohamed (Temple U), Jiaxing Qiu (Brookhaven National Lab)



Joint CTEQ Meeting and POETIC  
7th International Conference on  
Physics Opportunities at an Electron-Ion  
Collider  
Temple University  
November 14-18, 2016



**ECT\***  
EUROPEAN CENTRE FOR THEORETICAL STUDIES  
NUCLEAR PHYSICS AND RELATED AREAS




**The Proton Mass: At the Heart of Most Visible Matter**  
From Monday, 3 April 2017, 09:00 to Friday, 7 April 2017, 13:30

Speakers:  
NAC, Links, Safety




**EICUG MEETING - July 18-22, TRIESTE 2017**  
Hosting Institution: INFN, Sezione di Trieste  
in cooperation with Trieste University



**Programs & Workshops**

- 2017 Programs**
  - Toward Predictive Theories of Nuclear Reactions Across the Isotopic Chart (INT-17-1a) March 31, 2017
  - Microscopy of QGP Properties with Jets and Heavy Quarks (INT-17-1b) June 5, 2017
  - Double-beta Decay (INT-17-2a) July 14, 2017
  - Relativistic Signatures of R-process Nuclei (INT-17-2b) August 18, 2017
  - Spin and Momentum Tomography of Hadrons and Nuclei (INT-17-3) August 28 - September 29, 2017
- 2017 Workshops**
  - Probing QCD in Photon-Nucleus Interactions at RHIC and LHC: the Path to EIC (INT-17-65W) February 13 - 17, 2017
  - SIGN 2017: International Workshop on the Sign Problem in QCD and Beyond (INT-17-64W) March 20 - 24, 2017
  - Lattice QCD Input for Neutrinoless Double- $\beta$  Decay (INT-17-67W) July 6 - 7, 2017
  - The Flavor Structure of Nucleon Sea (INT-17-68W) October 2 - 13, 2017
  - Neutron-Antineutron Oscillations: Appearance, Disappearance, and Baryogenesis (INT-17-69W) October 23 - 27, 2017
- 2018 Programs**
  - Nuclear *ab-initio* Theories and Neutrino Physics (INT-18-1a) February 26 - March 30, 2018
  - Multi-Scale Problems Using Effective Field Theories (INT-18-1b) May 7 - June 1, 2018
  - Fundamental Physics with Electroweak Probes of Light Nuclei (INT-18-2a) June 12 - July 13, 2018
  - Advances in Monte Carlo Techniques for Many-Body Quantum Systems (INT-18-2b) July 30 - September 7, 2018
  - Probing Nucleons and Nuclei in High Energy Collisions (INT-18-3) October 1 - November 16, 2018

Programs related to EIC




relation to their physics  
in electroweak probe  
hadron production with  
in view of new discoveries of multi-quark





# Summary


- The EIC will profoundly impact our understanding of QCD and its dynamics  
→ the **structure of nucleons and nuclei** in terms of sea quarks & gluons
  - The EIC will enable **imaging** and provide **unprecedented kinematic reach** into **yet unexplored regions of phase spaces in QCD** with its high luminosity/energy, nuclei & beam polarization
- ⇒ High potential for discovery within QCD with broad impact beyond**
- Outstanding questions raised by the science at RHIC/LHC and CEBAF, have **naturally led to the EIC Science & design parameters**
  - **World wide interest** in collaborating on the EIC
  - Accelerator scientists at BNL and JLab together with international accelerator community is ready to provide the **intellectual and technical leadership for to realize the EIC**, a frontier accelerator facility.



# Summary

- The EIC will profoundly impact our understanding of QCD and its dynamics  
→ the **structure of nucleons and nuclei** in terms of sea quarks & gluons
  - The EIC will enable **imaging** and provide **unprecedented kinematic reach** into **yet unexplored regions of phase spaces in QCD** with its high luminosity/energy, nuclei & beam polarization
- ⇒ **High potential for discovery within QCD with broad impact beyond**
- Outstanding questions raised by the science at RHIC/LHC and CEBAF, have **naturally led to the EIC Science & design parameters**
  - **World wide interest** in collaborating on the EIC
  - Accelerator scientists at BNL and JLab together with international accelerator community is ready to provide the **intellectual and technical leadership for to realize the EIC**, a frontier accelerator facility.

**Future of nuclear science demands an Electron Ion Collider.  
The U.S. must lead the way.**



---

Thank you!

